

53883-183

4/17/2014

1/17



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

Control Solutions, Inc.
% Jane Miller
Biologic Inc.
115 Obtuse Hill
Brookfield, CT 06804

APR 17 2014

Subject: Application for Pesticide Notification (PRN 98-10)
T-Methyl G-Pro Granular
EPA Reg. No. 53883-183
Notification dated: March 14, 2014
EPA Decision Number 489482

Dear Ms Miller:

The Agency is in receipt of your Application for Pesticide Notification under Pesticide Registration Notice (PRN) 98-10. The Registration Division (RD) has conducted a review of this request for its applicability under PRN 98-10 and finds that the action(s) requested fall within the scope of PRN 98-10.

The label submitted with the application has been stamped "Notification" and will be placed in our records.

If you have any questions, you may contact Shaunta Hill at 703-347-8961 or via email at hill.shaunta@epa.gov.

Sincerely,

A handwritten signature in black ink that reads "Shajia B. Joyner".

Shajia B. Joyner
Product Manager 20
Fungicide Branch
Registration Division (7505P)
Office of Pesticide Programs



United States
Environmental Protection Agency
 Washington, DC 20460

<input type="checkbox"/>	Registration
<input type="checkbox"/>	Amendment
<input checked="" type="checkbox"/>	Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 53883-183	2. EPA Product Manager S. Joyner	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) T-Methyl G-Pro Granular	PM# 20	
5. Name and Address of Applicant (Include ZIP Code) Control Solutions, Inc. c/o Biologic, Inc. 115 Obtuse Hill Road Brookfield, CT 06804 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input checked="" type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

This is a notification to add marketing statements per PR Notice 98-10. This notification is consistent with the provisions of PR Notice 98-10 and the EPA regulations at 40 CFR 152.46 and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46 this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
				<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
* Certification must be submitted	If "Yes" Unit Packaging wgt.	No. per container	If "Yes" Package wgt	No. per container	Other (Specify) _____
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> on label	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled			<input type="checkbox"/> Other _____		

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Jane M. Miller	Title Agent	Telephone No. (Include Area Code) (203) 740-1200
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title Agent	
4. Typed Name Jane M. Miller	5. Date March 14, 2014	

March 14, 2014

Document Processing Desk (NOTIF)
Office of Pesticide Programs (7504P)
US Environmental Protection Agency
One Potomac Yard, 2777 S. Crystal Drive
Arlington, VA 22202-4501

Attention: Ms. Shaja B. Joyner (PM #20)

Re: "T-Methyl G-Pro Granular"; EPA Reg. No. 53883-183
Notification to Add Optional Marketing Statements

Dear Ms. Joyner:

On behalf of Control Solutions, Inc. please find enclosed this Application for Pesticide Notification for the above-mentioned product. This Notification is to add optional marketing statement to the front panel.

The proposed text will comply with PR-Notice 98-10 Section N. **Other Revisions - Minor label changes not described in Section II.A.-M. and Section III. which are related to FIFRA be made by notification, 3. provided they involve no change in the ingredients statement, signal word, use classification, precautionary statements, statements of practical treatment (First Aid), physical/chemical/biological properties, storage and disposal, or directions for use.**

This action fits the definition of a Notification and:


"This notification is consistent with the provisions of PR Notice 98-10 and the EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or the confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statement to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA."

The following documents are enclosed to process this Notification:

- 1. Application for Pesticide Notification (8570-1)
- 2. One (1) copy of the revised label with the additional text underlined/highlighted.
- 3. One (1) clean copy of the label.

Should you have any questions, or wish to reach me, please feel free to contact our office at 203-740-1200.

Sincerely,



Jane M. Miller
Agent to Control Solutions, Inc.

MASTER LABEL

EPA Approved Amended Label 12 18 2009
Notification to add Marketing Language 03 13 2014

4/17

T-Methyl G-Pro Granular
Systemic Fungicide for Turf and Ornamentals

For the prevention and control of turf diseases and the diseases of annual and perennial flowers, bedding plants, foliage plants, ground covers, plus deciduous and evergreen trees and shrubs.

[OPTIONAL MARKETING STATEMENTS:

- Covers [Treats up to] 5,000 sq. ft [10,000 sq. ft.]
Lawn disease control [Spring][Summer][Fall]
Also provides broad spectrum systemic control of soil borne diseases
Effective on 32 listed diseases
More than 25 diseases controlled! [see list on back of package]
May be used on cool & warm season grasses
Economical control of lawn turf diseases.
Use in combination with other fungicide to prevent the development of tolerant disease strains.
Your Prescription for a Picture Perfect Lawn™]

[ROOM FOR ARTWORK]

ACTIVE INGREDIENT

Thiophanate-methyl (dimethyl 4,4'-o-phenylenebis[3-thioallophanate])..... 2.08%

OTHER INGREDIENTS..... 97.92%

TOTAL:..... 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

Table with 2 columns: First Aid instructions (If in eyes, Swallowed, Inhaled, Skin/Clothing) and Hot Line Number information.

For spill or transportation emergencies call CHEMTREC (1-800-424-9300).

EPA Reg. No. 53883-183

EPA Est. No. XXXXX-XX-XXX

Control Solutions, Inc.
5903 Genoa-Red Bluff
Pasadena, TX 77507
Net Weight: ___ lbs.

NOTIFICATION

APR 17 2014

6/17

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific statements and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval (REI). The requirements in this box apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls over long-sleeved shirt and long pants,
- Chemical-resistant gloves made of any waterproof material,
- shoes plus socks
- Chemical-resistant headgear for overhead exposure

NON-AGRICULTURAL USE REQUIREMENT

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Turf and Landscape Uses: Do not enter or allow entry into treated areas until granules and dust have settled unless wearing personal protective equipment listed on this label.

COMMERCIAL APPLICATIONS

GENERAL INFORMATION

T-Methyl G-Pro Granular is a broad-spectrum fungicide for use on a wide variety of turf and ornamental diseases. T-Methyl G-Pro Granular may be applied using ground based, hand-operated (push) or mechanical (power) equipment; do not apply this product by hand or with hand-held rotary spreader (e.g., bellygrinder).

USE PRECAUTIONS

- This product may only be used by individuals/firms licensed or registered by the state to apply ornamental or turf pest control products.
- Do not apply this product through any type of irrigation system.
- Do not use on turf being grown for sale or other commercial use as sod.
- Do not graze animals on treated turf.
- Do not feed clippings to livestock or poultry.
- Use only properly calibrated application equipment when applying this product.
- Thoroughly clean all application equipment after each use.
- For application schedules appropriate to your area, contact your local State Extension Service for recommendations.

RESISTANCE MANAGEMENT

In order to prevent the development of tolerant strains of fungi, T-Methyl G-Pro Granular should only be used with other fungicides that use different modes of action. CONTROL SOLUTIONS, INC. does NOT recommend using T-Methyl G-Pro Granular with products containing the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can contribute to the development of disease tolerance. T-Methyl G-Pro Granular can remain useful for disease control as long as specified precautions are followed. A tolerant strain of fungi may be present if a T-Methyl G-Pro Granular treatment is found to be ineffective and you should promptly contact your State Agricultural Experiment Station or State Cooperative Extension Service for disease identification and advice on other suitable fungicide or disease control strategies.

MAXIMUM APPLICATION RATES AND MINIMUM RETREATMENT INTERVALS

Refer to the following table for the maximum treatment and seasonal application rates allowed based on the use site being applied to.

Use Site	Maximum Per-Treatment Application Rate of T-Methyl G-Pro Granular	Maximum Per-Season Application Rate of T-Methyl G-Pro Granular	Minimum Retreatment Interval
Residential or Public Areas [†]	131 lbs./acre (3 lbs./1000 sq. ft.)	523 lbs./acre (12 lbs./1000sq. ft.)	14-days
Golf Course Aprons, Greens and Tees	392 lbs./acre (9 lbs./1000 sq. ft.)	1048 lbs./acre (24 lbs./1000sq. ft.)	
Golf Course Fairways (except Florida)	262 lbs./acre (6 lbs./1000 sq. ft.)	262 lbs./acre (6 lbs./1000sq. ft.)	
Golf Course Fairways (Florida only) [‡]	131 lbs./acre (3 lbs./1000 sq. ft.)	131 lbs./acre (3 lbs./1000sq. ft.)	

[†] Maximum of four (4) applications per year.

[‡] Maximum of one (1) application per year.

COMMERCIAL TURF APPLICATIONS

T-Methyl G-Pro Granular may be used on all cool and warm season grasses such as Bentgrass, Bluegrass, Bermudagrass, Fescue, Ryegrass, St. Augustinegrass, Zoysiagrass, or their mixtures in the following sites:

- Commercial turf
- Residential turf
- Public turf (parks, athletic fields, schools, day care centers, etc.)
- Golf courses (greens, tees, fairways and aprons)


When used in accordance with the instructions in this label, T-Methyl G-Pro Granular is not phytotoxic to any of the above-mentioned grasses.

Application Instructions

- Apply under dry or moist leaf/soil conditions.
- For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10-0.25 inches of water.
- Treating aprons when treating golf greens is recommended.

Refer to the following table for specific rates and timing depending on the disease being treated. When treating under severe disease pressure, use the highest recommended rate listed below.

Disease(s)	Disease Specific Instructions	Application Rate of T-Methyl G-Pro Granular (Lbs./1000 sq. ft.)
<p>Dollar Spot <i>Moellerodiscus, Lanzia, Sclerotinia</i></p> <p>Large Brown Patch <i>Rhizoctonia solani</i></p> <p>Ascochyta Leaf Blight <i>Ascochyta spp.</i></p> <p>Copper Spot <i>Gloeocercospora sorghi</i></p> <p>Red Thread <i>Laetisaria fuciformis</i></p> <p>Fusarium Patch <i>Fusarium nivale</i></p> <p>Zoyzia Patch <i>Rhizoctonia solani</i></p>	<p>For Prevention: Apply prior to the expected development of the disease using the lowest rate listed.</p> <p>For Treatment: Apply when disease first appears and repeat applications every 14-days.</p> <p>NOTE: May be rotated with chlorothalonil, iprodione, or mancozeb (Protect™).</p>	<p>1.5-6</p>
<p>Pink Snow Mold <i>Microdochium nivale</i></p>	<p>Make first application prior to cessation of turf growth activity in middle-late Fall or early Winter, or in Spring during periods of thaw, or when disease symptoms first appear.</p> <p>Throughout the expected disease period, continue making applications at 14-day intervals.</p> <p>For best results, make the first application before the turf is completely dormant.</p>	<p>3-6</p>
<p>Anthracnose, foliar <i>Colletotrichum graminicola</i></p> <p>Anthracnose, basal <i>Colletotrichum graminicola</i></p>	<p>For Prevention: Once soil temperature reaches 60°F, make two applications at 14-day intervals.</p> <p>For Treatment: Apply at first sign of disease and continue applications at 14-day intervals.</p> <p>NOTE: May be rotated with chlorothalonil or triadimefon.</p>	<p>3-6</p> <p>6-9</p>



Disease(s)	Disease Specific Instructions	Application Rate of T-Methyl G-Pro Granular (Lbs./1000 sq. ft.)
Bermudagrass Decline <i>Gaeumannomyces graminis</i> var. <i>graminis</i> Take-All-Patch <i>Gaeumannomyces graminis</i> var. <i>aavenae</i>	Apply in mid-July or when disease symptoms first appear and repeat at 14-day intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic recommendations to maintain plant vigor.	
Cool Season Brown Patch <i>Rhizoctonia cerealis</i> Necrotic Ring Spot <i>Leptosphaeria korrea</i> Spring Dead Spot <i>Leptosphaeria korrea</i>	For Prevention: Apply in the fall prior to cessation of turf growth. Make a second application in early Spring when disease first appears or when soil temperatures reach 55-60°F, whichever occurs first. For Treatment: Apply in early Spring when disease first appears and repeat applications every 14-days.	
Coprinus Snow Mold <i>Coprinus psychromorbidus</i>	Make two treatments at 21-day intervals in late fall to early winter. Make the last application just prior to first permanent snow cover.	
Fusarium Blight <i>Fusarium roseum</i> and <i>F. triticum</i>	Apply when disease first appears. Make repeat applications at 14-day intervals.	6-9
Gray Leaf Spot (Blast) <i>Pyricularia grisea</i>	Make a preventative application prior to the expected onset of disease development. Continue applications at 14-day intervals as needed.	
Leaf Spot <i>Drechslera</i> spp. Leaf, Crown, and Root Diseases <i>Bipolaris</i> , <i>Curvularia</i> , and <i>Exserohilum</i>	Apply when disease first appears and make applications at 14-day intervals as needed. NOTE: For severe cases, rotations with chlorothalonil, iprodione, or mancozeb (Protect™) are recommended.	
Stripe Smut <i>Ustilago striiformis</i>	For Prevention: Apply in spring and fall. For Treatment: Make two applications, the first when the disease first appears and the second 14-days later.	
Summer Patch <i>Magnaporthe poae</i>	For Prevention: Using a 21-day reapplication interval, make three applications starting in late April or early May. For Treatment: Apply at 14-day intervals when disease first appears. Rotations may be used as part of the three-application program.	

Disease(s)	Disease Specific Instructions	Application Rate of T-Methyl G-Pro Granular (Lbs./1000 sq. ft.)
Bentgrass Dead Spot <i>Ophiosphaerellia agrostis</i>	Apply in early June or when recommended by your local Agricultural Extension Service. Reapply at 14-day intervals as needed. For season- long disease prevention, rotations and/or tank mix combinations may be used.	6-9

COMMERCIAL HORTICULTURAL APPLICATIONS

T-Methyl G-Pro Granular is a broad spectrum systemic fungicide for control of soil borne diseases effecting bulbs, corms, rhizomes, roots, crowns, and stems caused by:

- *Botrytis*
- *Gliocladium*
- *Ramularia*
- *Thielaviopsis*
- *Cylindrocladium*
- *Myrothecium*
- *Rhizoctonia*
- *Fusarium*
- *Penicillium*
- *Sclerotinia*

T-Methyl G-Pro Granular may be used on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs found in Nursery, Greenhouse, Interiorscape, and Landscape sites.

When using this product for commercial horticultural applications, be sure to follow the guidelines in the General Information Section above.

NOTE: Because T-Methyl G-Pro Granular does not control *Phthium* or *Phytophthora*, rotating with metalaxyl, mefenoxam, etridiazole, propamocarb, mono and dipotassium salts of phosphorous acid or fosetyl-AI is required to control *Pythium* and *Phytophthora*.

USE PRECAUTIONS

- This product is not recommended for use on Swedish Ivy (*Nephrolepis exhalta*), Boston Fern (*Plectranthus australis*), and Easter Cactus (*Hatiora gaertneri*).
- Do not apply to home orchards / fruit trees after fruit set.
- Do not use fruit or nuts from treated trees for food or feed.
- For ground applications, do not exceed 300 pounds active ingredient from ALL thiophanate-methyl containing products per acre per crop season.
- For cut flowers, the maximum application rate is 0.5 pounds active ingredient per acre per application.
- Because it is impossible to test this product on all species and cultivars, a preliminary trial on a small scale is recommended before a full treatment is applied to any plant type not shown on this label. When conducting the trial, wait 5-7 days after treatment to evaluate results.

Application Instructions

T-Methyl G-Pro Granular may be applied as a surface, fly-on, broadcast or over-the-top application or media incorporated.

Surface and Broadcast Applications to established plants:

- Apply 6 lbs. of T-Methyl G-Pro Granular / 1000 sq. ft.
- Apply under dry or moist leaf / soil conditions.

- For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10-0.25 inches of water.

Media Incorporation:

- Mix at a rate of 2-3 lb/cubic yards before seeding or transplanting of desired crop.
- If not activated by the application of moisture, media incorporated with T-Methyl G-Pro Granular may be stockpiled until needed for later use.

For Preventive Control:

- Prior to planting, apply as a media incorporation using the rate listed above.
- After seeding or transplant to propagation beds, containers, pots, or nursery or landscape beds, apply as a broadcast, fly-on, surface or over the top application using the Surface and Broadcast Applications instructions above.

For Treatment:

- When disease first appears after seeding or transplanting, apply using the Surface and Broadcast Applications instructions above.
- To maintain protection, make repeat applications every 21-28 days.


STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Pesticide Storage: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.

Pesticide Disposal: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spilled bait, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

Container Disposal: Non-refillable container. Do not reuse or refill this container. Empty container through use according to label instructions. Offer for recycling, if available. If recycling is not available; then dispose of container in a sanitary landfill or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.



[HOMEOWNER USE LABEL LANGUAGE]

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION - Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt, long pants, and shoes with socks.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or areas where surface water is present or to intertidal areas below the mean high water mark. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will directly contact any person or pet.

Keep people and pets out of the area during application. Do not enter or allow others to enter the treated areas until dusts have settled.

HOMEOWNER TURF AND ORNAMENTAL APPLICATIONS

GENERAL INFORMATION

T-Methyl G-Pro Granular is a broad-spectrum fungicide for use on a wide variety of turf diseases. T-Methyl G-Pro Granular may be applied using ground based, hand-operated (push) or mechanical (power) equipment; do not apply this product by hand or with hand-held rotary spreader (e.g., bellygrinder). If a rate range is given, use the higher rate under conditions of severe disease pressure. Contact your State Extension service for recommendations for application schedules appropriate to your area.


USE PRECAUTIONS

- Do not apply this product through any type of irrigation system.
- Do not apply by hand or with handheld rotary spreader (e.g. belly grinder).
- Do not graze animals on treated turf.
- Do not feed clippings to livestock or poultry.
- Use only properly calibrated application equipment when applying this product.
- Thoroughly clean all application equipment after each use.
- This product is not guaranteed to be compatible with any other pesticides.

RESISTANCE MANAGEMENT

In order to prevent the development of tolerant strains of fungi, T-Methyl G-Pro Granular should only be used with other fungicides that use different modes of action. CONTROL SOLUTIONS, INC. does NOT recommend using T-Methyl G-Pro Granular with products containing the active ingredient thiabendazole because thiabendazole has a similar chemistry and mode of action and can contribute to the development of disease tolerance. T-Methyl G-Pro Granular can remain useful for disease control as long as specified precautions are followed. A tolerant strain of fungi may be present if a T-Methyl G-Pro Granular treatment is found to be ineffective and you should promptly contact your State Agricultural

Disease(s)	Disease Specific Instructions	Application Rate (Lbs./1000 sq. ft.)	
<p>Dollar Spot <i>Moellerodiscus, Lanzia, Sclerotinia</i></p> <p>Large Brown Patch <i>Rhizoctonia solani</i></p> <p>Ascochyta Leaf Blight <i>Ascochyta spp.</i></p> <p>Copper Spot <i>Gloeocercospora sorghi</i></p> <p>Red Thread <i>Laetisaria fuciformis</i></p> <p>Fusarium Patch <i>Fusarium nivale</i></p> <p>Zoyzia Patch <i>Rhizoctonia solani</i></p>	<p>For Prevention: Apply prior to the expected development of the disease using the lowest rate listed.</p> <p>For Treatment: Apply when disease first appears and repeat applications every 14-days.</p> <p>NOTE: May be rotated with chlorothalonil, iprodione, or mancozeb (Protect™).</p>	<p>1.5-3</p>	
<p>Pink Snow Mold <i>Microdochium nivale</i></p>	<p>Make first application prior to cessation of turf growth activity in middle-late Fall or early Winter, or in Spring during periods of thaw, or when disease symptoms first appear.</p> <p>Throughout the expected disease period, continue making applications at 14-day intervals.</p> <p>For best results, make the first application before the turf is completely dormant.</p>		
<p>Anthracnose, foliar <i>Colletotrichum graminicola</i></p> <p>Anthracnose, basal <i>Colletotrichum graminicola</i></p>	<p>For Prevention: Once soil temperature reaches 60°F, make two applications at 14-day intervals.</p> <p>For Treatment: Apply at first sign of disease and continue applications at 14-day intervals.</p> <p>NOTE: May be rotated with chlorothalonil or triadimefon.</p>		



Disease(s)	Disease Specific Instructions	Application Rate (Lbs./1000 sq. ft.)
Bermudagrass Decline <i>Gaeumannomyces graminis</i> <i>var. graminis</i> Take-All-Patch <i>Gaeumannomyces graminis</i> <i>var. aavenae</i>	Apply in mid-July or when disease symptoms first appear and repeat at 14-day intervals for suppression. Use higher rates under most severe disease expression. Follow proper agronomic instructions to maintain plant vigor.	
Cool Season Brown Patch <i>Rhizoctonia cerealis</i> Necrotic Ring Spot <i>Leptosphaeria korrea</i> Spring Dead Spot <i>Leptosphaeria korrea</i>	For Prevention: Apply in the fall prior to cessation of turf growth. Make a second application in early Spring when disease first appears or when soil temperatures reach 55-60°F, whichever occurs first. For Treatment: Apply in early Spring when disease first appears and repeat applications every 14-days.	
Coprinus Snow Mold <i>Coprinus psychromorbidus</i>	Make two treatments at 21-day intervals in late fall to early winter. Make the last application just prior to first permanent snow cover.	
Fusarium Blight <i>Fusarium roseum</i> and <i>F. triticum</i>	Apply when disease first appears. Make repeat applications at 14-day intervals.	1.5-3
Gray Leaf Spot (Blast) <i>Pyricularia grisea</i>	Make a preventative application prior to the expected onset of disease development. Continue applications at 14-day intervals as needed.	
Leaf Spot <i>Drechslera spp.</i> Leaf, Crown, and Root Diseases <i>Bipolaris, Curvularia, and Exserohilum</i>	Apply when disease first appears and make applications at 14-day intervals as needed. NOTE: For severe cases, rotations with chlorothalonil, iprodione, or mancozeb (Protect™) are recommended.	
Stripe Smut <i>Ustilago striiformis</i>	For Prevention: Apply in spring and fall. For Treatment: Make two applications, the first when the disease first appears and the second 14-days later.	
Summer Patch <i>Magnaporthe poae</i>	For Prevention: Using a 21-day reapplication interval, make three applications starting in late April or early May. For Treatment: Apply at 14-day intervals when disease first appears. Rotations may be used as part of the three-application program.	

16/17

HOMEOWNER HORTICULTURAL APPLICATIONS

T-Methyl G-Pro Granular is a broad spectrum systemic fungicide for control of soil borne diseases effecting bulbs, corms, rhizomes, roots, crowns, and stems caused by:

- *Botrytis*
- *Gliocladium*
- *Ramularia*
- *Thielaviopsis*
- *Cylindrocladium*
- *Myrothecium*
- *Rhizoctonia*
- *Fusarium*
- *Penicillium*
- *Sclerotinia*

T-Methyl G-Pro Granular may be used on annual and perennial flowers, bedding plants, foliage plants, ground covers, and deciduous and evergreen trees and shrubs.

When using this product be sure to follow the guidelines in the General Information Section above.

NOTE: Because T-Methyl G-Pro Granular does not control *Phthium* or *Phytophthora*, rotating with metalaxyl, mefenoxam, etridiazole, propamocarb, mono and dipotassium salts of phosphorous acid or fosetyl-AI is required to control *Pythium* and *Phytophthora*.

USE PRECAUTIONS

- This product is not recommended for use on Swedish Ivy (*Nephrolepis exhalta*), Boston Fern (*Plectranthus australis*), and Easter Cactus (*Hatiora gaertneri*).
- DO NOT apply to home orchards / fruit trees.
- For cut flowers, the maximum application rate is 0.5 pounds active ingredient per acre per application.
- Because it is impossible to test this product on all species and cultivars, a preliminary trial on a small scale is recommended before a full treatment is applied to any plant type not shown on this label. When conducting the trial, wait 5-7 days after treatment to evaluate results.

Application Instructions

T-Methyl G-Pro Granular may be applied as a surface, fly-on, broadcast or over-the-top application or media incorporated.

Surface and Broadcast Applications to established plants:

- Apply 3 lbs. of T-Methyl G-Pro Granular / 1000 sq. ft.
- Apply under dry or moist leaf/soil conditions.
- For best results, activate the application within 24 hours with irrigation or natural rainfall. If activating by irrigation, apply 0.10-0.25 inches of water.

Media Incorporation:

- Mix at a rate of 2-3 lb/cubic yards before seeding or transplanting of desired crop.
- If not activated by the application of moisture, media incorporated with T-Methyl G-Pro Granular may be stockpiled until needed for later use.

For Preventive Control:

- Prior to planting, apply as a media incorporation using the rate listed above.
- After seeding or transplant to propagation beds, containers, pots, or nursery or landscape beds, apply as a broadcast, fly-on, surface or over the top application using the Surface and Broadcast Applications instructions above.

For Treatment:

- When disease first appears after seeding or transplanting, apply using the Surface and Broadcast Applications instructions above.
- To maintain protection, make repeat applications every 21-28 days.

SPRADER SETTINGS (PER 1,000 SQ. FT.)		
	1.5 lbs. (Prevention) SETTING	3.0 lbs. (Treatment) SETTING
DROP SPREADERS:		
Republic EZ #530	1.5	2.5
Scotts AccuGreen	3	5
Sears Drop	3.5	5
ROTARY SPREADERS:		
Earthway	7	9
Republic EZ #540	1.5	3
Scotts SpeedyGreen	3.5	6.5
Sears Rotary	1.5	3

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.
Pesticide Storage: Store in original container in a dry place. Store in locked area out of reach of children and domestic animals.
Pesticide Disposal: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WARRANTY STATEMENT

The following limitations of warranty shall apply to the extent that they are consistent with applicable law. Control Solutions, Inc. warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. This product is not guaranteed to be compatible with any other pesticides. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Control Solutions, Inc. To the extent consistent with applicable law, Control Solutions, Inc. shall in no event be liable for consequential, special, or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. Except, as expressly provided herein, Control Solutions, Inc. makes no warranties, guarantees, or representations of any kind, either expressed or implied, or by usage of trade, statutory or otherwise, with regard to the product sold, including, but not limited to merchantability, fitness for a particular purpose, use or eligibility of the product for any particular trade usage. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damage resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall be damages not exceeding the purchase price paid for this product or, at Control Solutions, Inc. election, the replacement of this product.

